GENERAL INDEX TO VOLUME XXXIV

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italics;* and previously published names and all other matter, in ordinary type.

A

Adiantum pedatum, 122
Adlumia, 191; fungosa, 188, 239
Alkaloidal properties of Corydalis, 195
American origin of the cultivated Cucurbits, 101
Anderson, Edgar: Field studies of Guatemalan maize, 433; William L. Brown and. The northern flint corns, 1
Andrews, Henry N.: John Henry Britts—physician and fossil hunter, 115; and Ellen M. Kern. The Idaho Tempskyas and associated fossil plants, 119; and Lee W. Lenz. Fossil Polypores from Idaho, 113
Anemia elegans, 149; Fremonti, 152
Antigua, Guatemala, typical ear of corn from region about, *452*, *456*
Archeological maize in the United States, 8, 10, 13, *28*
Asclepias incarnata, 356; speciosa × syriaca, 358; syriaca, 355, 358; tuberosa, 353, cl. interior, *353*, ssp. interior, 353, 361, intraspecific differentiation of, 409, ssp. Rolfsii, 361, ssp. tuberosa, 361
Asclepias tuberosa, Some dynamics of leaf variation in, 353
Asclepias tuberosa: biology of, 354; cytogenetics of, *358*, 359; ecology of, 359; hybridization in, 391; leaf types of three subspecies, 361, *366*, *367;* phenocontour maps, 370; population studies, 362; reproductive habit, 355; roadside profile of, 394, *395*, *401;* compared with herbarium profile, 403, *407;* vegetative habit, 354
Austroclepis australis, 146

B

Bennettiales, fossil, 153
Bolivia, wild rubber in, 266
Borckhausenia, 197
Brettanomyces, 33
Britts, John Henry—physician and fossil hunter, 115, *115*
Brown, William L., and Edgar Anderson. The northern flint corns, 1
Bulbocapnos, 197, 207
Burkholder's medium, growth of yeast cultures on, *76*, 81, 85, 95
Butterflyweed, see Asclepias tuberosa

C

Candida Guilliermondii, 33; *mortifera*, 33
Caoutchoua elastica, 292
Capnites, 197, 207
Capnodes, 197; *aureum*, 229; *Bidwellianum*, 201; *Caseanum*, 201; *crystallinum*, 217; *flavulum*, 215; *pauciflorum*, 207; *Scouleri*, 199
Capnoides, 188, 197, 209; *brachycarpum*, 204; *Brandegei*, 203; *campestre*, 222; *curvisiliquum*, 223, 226; *Cusickii*, 205; *Engelmannii*, 230; *euchlamydeum*, 230; *glauca*, 211; *Halei*, 217, 222, 223; *hastatum*, 206; *macrorrhiza*, 230; *micranthum*, 219; *montanum*, 234; *pachylobum*, 234; *pauciflorum*, 207; *Scouleri*, 199; *sempervirens*, 188, 211; *Wetherillii*, 230
Carnation (Dianthus caryophyllus), Inheritance in the, III. Inheritance of flower color, 39, *72*, *74*
Cells, relation between viability of, and inability to grow on deficient medium, 83
Ceratopteris thalictroides, 149
Chimaltenango, Guatemala, corn from, *462*
Chromosomal complements of Corydalis, 194
Chromosome knob numbers in dent corn inbreds, *5*
Cibotium sp., compared with Tempskya Wessellii, 133
Cisticapnos, 188
Clepsydropsis australis, 146
Clons of butterflyweed, *353*, 354
Coal mines, fossil plants of, collected by Dr. Britts, 115
Color in carnations, inheritance of, *39*, *72*, *74;* acyanic group, 41; cyanic group, 48; transition group, 56; variegated group, 57
Coniferales, fossil, 153
Corn: archeology of, 10; breeding, 16, 449; brewing varieties, 445; dent varieties, 1, 15; field studies of, in Guatemala, 433; hybridization of, 15; Mexican, 14; "mul-

tiplication" in Guatemalan varieties, 448; northern flint, 1; of the Corn Belt, 14; popping varieties, 445; prehistoric collections, 9, *28*

Corydalis: Monograph of the North American species of, 187; morphology of, 188, of flowers, *252–258*; taxonomy, 197

Corydalis, 187; *Albertae*, 229; *Allenii*, 199; *annua*, 211; aurea ssp. **aurea**, 229, ssp. **occidentalis**, 234; *aurea* var. *australis*, 222, *β crystallina*, 217, var. *curvisiliqua*, 226, *α flavula*, 215, var. *macrantha*, 229, var. *micrantha*, 219, var. *occidentalis*, 234, var. *parviflora*, 229, var. *robusta*, 229, var. *typica*, 229; *biaurita*, 239; *Bidwelliae*, 201; *bilimbata*, 234; *brachycarpa*, 204; *bracteosa*, 239; *Brandegei*, 203; bulbosa, 194; *campestris*, 222; *canadensis*, 239; Caseana, 201, ssp. **brachycarpa**, 204, ssp. **Brandegei**, 203, ssp. **Caseana**, 201, *256*, ssp. **Cusickii**, 205, ssp. **hastata**, 206; cava, 194; *chihuahuana*, 234; *crassipedicellata*, 234; crystallina, 217, var. *strictissima*, 217; *Cucullaria*, 239; curvisiliqua, 226, ssp. **curvisiliqua**, 226, ssp. **grandibracteata**, 227; *curvisiliqua* var. *grandibracteata*, 228, var. *tenerior*, 222; *curvisiliquaeformis*, 234; *Cusickii*, 205, var. *hastata*, 206; *densicoma*, 230; *Engelmannii*, 230, var. *exaltata*, 230; *euchlamydea*, 230; sect. Eucorydalis, 209; *eximia*, 239; *flavidula*, 215; flavula, 215; *formosa*, 239; *fungosa*, 239; *Geyeri*, 215; *glauca*, 211; *Goodd*ingii, 229; *Halei*, 222; *hastata*, 206; *Hendersonii*, 205; *hypecoiformis*, 229; *idahoensis*, 205; *isopyroides*, 230, var. *Mearnsii*, 230; *Jonesii*, 229, var. *stenophylla*, 234; lutea, 239; *macrophylla*, 199; *macrorrhiza*, 229; micrantha, 219, ssp. **australis**, 222, ssp. **micrantha**, 219, ssp. **texensis**, 225; *micrantha* var. *diffusa*, 219, var. *leptosiliqua*, 222, var. *pachysiliquosa*, 219; *monilifera*, 229, var. *ferruginifera*, 219; *montana*, 234; ochotensis, 239; *oregana*, 230; *pachyloba*, 234; paeoniaefolia, 239; pauciflora, 207, *258*, var. *Chamissonis*, 207, var. *parviflora*, 207; sect. Pes-gallinaceus, 207; pseudomicrantha, 237, var. *Griffithsii*, 234; pumila, 194; sect. **Ramoso-sibiricae**, 198; *rosea*, 211; Scouleri, 199, 239, *256*; sempervirens, 211; *tenuifolia*, 239; *tortisiliqua*, 229, var. *longibracteata*, 230; *washingtoniana*, 229; *Wetherillii*, 229; *wyomingensis*, 229, var. *lativaginata*, 230

Corydallis, 197

Cryptoceras, 197

Cucumber, "Small Sugar," 102, 108

Cucumer: Meer, 102, *108*; Türckisch, 102, *108*

Cucurbita, 101; capitata, 102; digitata, 103; ficifolia, 101, 104, 106; indica, 102, *111*; Lagenaria, 106; maxima, 101, 103, 107, *111*; melanosperma, 103, 104; mixta, 106; moschata, 101, 102, 104, 105, 106; Pepo, 101, *108*, *111*, varieties of, 102, 106, *111*; perennis, 103; verrucosa, 102, 106, *111*

Cucurbits: cultivated, American origin of the, 101; evidence from the herbals, 101; evidence for the New World origin, 105; recent botanical evidence, 106; survey of old and recent botanical evidence, 103

Cuidad Vieja, Guatemala, corn from, *457*

Cupressinoxylon sp., 153, transverse and tangential section of, associated with Tempskya, *182*

Cushaw, 105, 106

Cycad species, compared with Tempskya Wessellii, 133

Cycadeoidea, 151, 153

Cysticapnos, 188

D

Dianthus caryophyllus, Inheritance of flower color in, 39

Dicentra, 191; canadensis, 239; Cucullaria, 239; eximia, 239; formosa, 239

Dicksonia fibrosa, 131, 142, 145

Dicotyledoneae, fossil, some notes and methods on, 153

Dodge, Carroll W. Mycocandida riboflavina, 31

Dueñas, Guatemala, corn from, *455*, *456*, *458*

Dynamics of leaf variation in Asclepias tuberosa, Some, 353

E

Economic aspects of Hevea, in Peru, 261

Economic importance of species of Corydalis, 195

Ethnobotany of Hevea, 303, of maize, 434

F

Fermentation of galactose and maltose in Saccharomyces, 95

Ferns, fossil, 132

Flower color in carnation, inheritance of, 39; color groups, 41
Fomes, 113; applanatus, 114; idahoensis, fossil, 113, *114;* pinicola, 114
Fossil hunter, John Henry Britts, physician and, 115
Fossil plants: The Idaho Tempskyas and associated, 119; collected by Dr. John Henry Britts, 115; collected by C. Henry Thomas, 185
Fossil polypores from Idaho, 113
Fumaria, 188; *aurea*, 229; *flavula*, 215; *glauca*, 211; *pauciflora*, 207; *sempervirens*, 211; *vesicaria*, 188

G

Galactose, ability of yeast cultures to ferment, 96
Geissman, T. A., Gustav A. L. Mehlquist and. Inheritance in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39
Genes, organized, versus molecular, 92
Geology of South America, 299
Gleichenites coloradensis, 152
Gourds, 106
Guatemalan maize: Field studies of, 433, *439*, *440;* Appendix, *453–467;* brewing varieties, 445; common varieties, *452–467;* compared with Mexican maize, 438; measurements of collections, 437; popping varieties, 445; salpor, 448

H

Hemitelia crenulata, 141, 144, *178;* Smithii, 143, *180*
Herbals, evidence from, of American origin of Cucurbita, 101
Herbarium: use of, for plant population studies, 362; profile compared with roadside profile, 403, *407*
Hevea, A study of (with its economic aspects) in the Republic of Peru, 261; cytology of, 287; distribution of, in Peru, 309, map showing, *310;* ethnobotany of, 303; hybridization of, 286, 305; morphology of, 264, buds and flowers, 280, *330*, *334–336*, leaflets, 277, short-shoots, 270, *328–332;* taxonomy, 292; variation, 287.
Hevea, 292; Benthamiana, 276, 304, *336*, *344–348;* brasiliensis, 276, 305, *328*, *332*, *340*, *344–348*, var. *acreana*, 306, var. *angustifolia*, 305, var. *cuneata*, 294, mut. Granthami, 266, × guianensis var. lutea, 314, var. *janeirensis*, 306, var. *latifolia*, 305, × pauciflora, 317, *352*, f. *Randiana*, 306, var. *Randiana*, 306, var. *stylosa*, 305, f. *subconcolor*, 306, f. *typica*, 306; camporum, 265; *caucho*, 292; *collina*, 292; *confusa*, 300; *cuneata*, 294; *Foxii*, 319; *glabrescens*, 319, 320; guianensis, 276, 292, *350*, *334–348*, var. *collina*, 292, var. *cuneata*, 292, 294, var. lutea, 293, *334–348*, var. marginata, 319, ssp. *occidentalis*, 292; var. occidentalis, 292, × pauciflora, 318, ssp. *typica*, 292; guianensis var. lutea, 293, × Benthamiana, 319, × pauciflora, 320, *352*, × pauciflora × brasiliensis, 321, f. *peruviana*, 294; *humilior*, 321; *janeirensis*, 305; *Kunthiana*, 305; *lutea*, 293, var. *cuneata*, 294, f. *pilosula*, 294, var. *pilosula*, 294, var. *typica*, 294; *membranacea*, 300, f. *leiogyne*, 300, var. *leiogyne*, 300; microphylla, 276, 285, 292, *342–350;* *minor*, 265; *nigra*, 292; nitida, 276, 297, *338*, *344–350*, var. toxicodendroides, 265; *paludosa*, 321; pauciflora, 276, 300, *328*, *338*, *344–348*, ssp. *coriacea*, 300, var. *coriacea*, 300, ssp. *typica*, 300; *peruviana*, 293; *Randiana*, 305; rigidifolia, 276, 291, *332*, *340*, *344–350;* *Sieberi*, 305; Spruceana, 276, 285, 291, *330*, *336*, *344–348;* *viridis*, 297, var. *toxicodendroides*, 265
Hybrid yeasts, 75
Hybridization: in Asclepias tuberosa, 379; in Hevea, 286
Hybrids, putative, of Hevea in Peru, 313

I

Idaho: Fossil polypores from, 113; northeast corner of Lanes Creek quadrangle, 122, *123;* Tempskyas and associated fossil plants from, 119; Some comments on the discovery of Tempskya in the vicinity of Wayan, by C. Henry Thomas, 185; view of hills southeast of Wayan, *158*
Inheritance: in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39; Mendelian, of genes affecting vitamin-synthesizing ability in Saccharomyces, 95
Insect visitants of Asclepias tuberosa, 356, *356*

J

Jatropha elastica, 292

K

Kern, Ellen M., Henry N. Andrews and. The Idaho Tempskyas and associated fossil plants, 119
Kloeckera (Pseudosaccharomyces), 33

L

Lagenaria, 105; siceraria, 101
Leaf variation in Asclepias tuberosa, 353
Lenz, Lee W., Henry N. Andrews and. Fossil polypores from Idaho, 113
Lindegren, Carl C.: and Gertrude Lindegren. Mendelian inheritance of genes affecting vitamin-synthesizing ability in Saccharomyces, 95; and Caroline Raut. A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing "mutants" in yeasts, 85; The effect of the medium on apparent vitamin-synthesizing ability in Saccharomyces, 75
Lindegren, Gertrude, Carl C. Lindegren and. Mendelian inheritance of genes affecting vitamin-synthesizing deficiencies of microorganisms, 95
Lorain, John, descriptions of early corn varieties by, 15

M

Macock, 105
Maíz chapolote, 44; maíz huajaqueña, 458; maíz negrito, 459
Maize, Guatemalan: Field studies of, 433; see Corn and Zea Mays
Maltose, ability of yeast cultures to ferment, 96
Mass collection studies of Asclepias tuberosa, 363
Media: Burkholder's, 75, growth of yeast cultures on, 76, 81, 85, 95; formulas of different synthetic, 78; Hutner's, 75, growth of yeast cultures on, 76, 82
Medio Monte, Guatemala, corn from, *456*
Medium, The effect of, on apparent vitamin-synthesizing deficiencies of microorganisms, 75
Mehlquist, Gustav A. L., and T. A. Geissman. Inheritance in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39
Melibiose-fermenting capacity of various yeast cultures, 77
Melons, 104
Melo-pepo, 102; clypeatus, 102; compressus, 102; latior Clypeiformis, 102, *111*; teres, 102
Mendelian inheritance of genes affecting vitamin-synthesizing ability in Saccharomyces, 95
Microorganisms, The effect of the medium on apparent vitamin-synthesizing deficiencies of, 75
Milkweed, see Asclepias
Missouri, paleobotanical work in, 115
Mixco, Guatemala, corn from, *464*
Mutants in yeasts, pantothenate-synthesizing, A direct relationship between pantothenate concentration and the time required to induce the production of, 85
Mylocyprinus robustus, 113
Mycocandida riboflavina, 31, *34*, *36*

N

Neckeria, 188, 197; *aurea*, 229, var. *occidentalis*, 234; *curvisiliqua*, 226; *flavula*, 215; *glauca*, 211; *micrantha*, 219; *sempervirens*, 211
Northern flint corns, 1: chromosome knob numbers for, 6, 10, 17; morphology of, 2; origin of, 18; varieties from the northeast, 6, from the Great Plains and Midwest, 7; varieties of: Dakota White, *22*, *pl. 3*, Longfellow Flint, *3*, *12*, Parker's Flint, *22*, *pl. 2*, Spanish Popcorn, *22*, *pl. 5*, Stevens' Flint, *11*, *22*, *pl. 1*, Twelverow Dakota, *22*, *pl. 4*.

O

Odoptera, 197; *aurea*, 229
Ownbey, Gerald B. Monograph of the North American species of Corydalis, 187

P

Paleobotanical collecting in Missouri, 115, in Idaho, 113, 119
Paleogeography of eastern North America, 376
Pantothenate: concentration, A direct relationship between, and the time required to induce the production of pantothenate-synthesizing mutants in yeasts, 85; genes affecting synthesis of, 95
Paraminobenzoic acid, genes affecting synthesis of, 97

Paramos, Guatemala, corn from, *463*
Patzún, Guatemala, corn from, *439*, *452*
Pepo, 103; Indicus minor angulosus, 103, *111*; maximus indicus compressus, 102, *111*; maximus oblongus, 102; minor clypeatus, 103, minor oblongus, 103; minor rotundus, 103
Pepones lati, 102, *111*
Peru: A study of Heavea in the Republic of, 261; map of, *310*
Phenocontour maps, use of, in study of Asclepias tuberosa, 370
Phoenix dactilifera, 132
Pioneer Hi-Bred Corn Co., experimental plots of, 1
Pistolochia, 197
Polypores, fossil, from Idaho, 113
Popcorn in Guatemala, 454, balls of, 447, being sold in market-place, *439*
Population patterns of three subspecies of Asclepias tuberosa, 362
Prehistoric corn collections, map of eastern United States showing, *13*
Pseudofumaria, 188
Pseudo-Fumaria, 197
Pumpkin, field, 102
Pyridoxine, genes affecting synthesis of, 97

Q

Quezeltenango, Guatemala, corn from, *452*, *453*, *466*, *467*

R

Raut, Caroline, Carl C. Lindegren and. A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing mutants in yeasts, 85; The effect of the medium on the apparent vitamin-synthesizing deficiencies of microorganisms, 75
Roadside profile of butterflyweed, analysis of, 394, compared with herbarium profile, 403, *407*
Rubber, Brazilian, 266; economic importance of, 261; history of, in South America, 261

S

Saccharomyces: Mendelian inheritance of genes affecting vitamin-synthesizing ability in, 95; carlsbergensis, effect of medium on vitamin synthesis with, 75, 91, pedigree of a hybrid between, and S. cerevisiae, 79; cerevisiae, effect of medium on vitamin synthesis with, 75, 85, hybrid of, heterozygous for mating type and fermentative ability, 96; *fragilis*, 31, 34
Salcajá, Guatemala, corn from, *454*
Salpor in Guatemala, 448
San Antonio Aguascalientes, Guatemala, corn from, *439*, *455*
San Juan de l'Obispo, Guatemala, corn from, *460*
San Lucas, Guatemala, corn from, *461*
San Miguel Morazón, Guatemala, corn from, *462*
Santa Lucía, Guatemala, corn from, *459*
Seibert, R. J. A study of Hevea (with its economic aspects) in the Republic of Peru, 261
Siphonia apiculata, 293; *brasiliensis*, 305; *brevifolia*, 293; *Cahuchu*, 292; *elastica*, 292; *guianensis*, 292; *Kunthiana*, 305; *lutea*, 293; *nitida*, 297; *pauciflora*, 300; *ridleyana*, 306
Sophorocapnos, 197
Squash, 101: "Acorn," 103; "Golden Custard," 102, *111*; "Perfect Gem," 103; "Table Queen," 103, *111*; Warted, 106; "White Bush Scallop," 102, *111*
Syringospora, 34

T

Tassel branches in northern flint corns, 2, *4*
Tempskya, 119; grandis, 146, 147; Knowltoni, 121, 147; minor, 130, 147; rossica, 146, 147; Schimperi, 149; Wessellii, 124, compared with living plants, 133; Whitei, 124; wyomingensis, 124, 147, 151
Tempskya: annuli spores and fragments, *144*; associated plant remains, 151; as animal food, 152; compared with other plants, 143; disc specimens, 127, *166*, *174*; longitudinal sections through, *138*; restoration showing probable habit, *140*; roots, 139, *168*, *170*; spores and sporangia, 149; stems, *136*, *172–176*; taxonomic considerations, 146; transverse sections of specimens, *126*, *135*, *137*, *176*; trunks, 125, 131, *160–164*
Tempskyas, The Idaho, and associated fossil plants, 119
Thiamin, genes affecting synthesis of, 95
Thomas, Mr. C. Henry: fossil-collecting activities of, 120; with a portion of his Tempskya collection, *158*; Some comments on the discovery of Tempskya in the vicinity of Wayan, Idaho, by, 185
Thyrsopteris elegans, 149

Todea barbara, 145
Tripsacum, 443; australe, 19

V

Variation: leaf, in Asclepias tuberosa, 353; pattern of, in Zea Mays, 433
Vegetable marrow, 102
Viability of cells, relation between, and inability to grow on deficient medium, 83
Vitamin-synthesizing: ability in Saccharomyces, Mendelian inheritance of, 95; deficiencies of microorganisms, The effect of the medium on apparent, 75
Vitamins, effect of multiple deficiencies of, on growth of Saccharomyces cerevisiae, 82

W

Whitaker, Thomas W.: American origin of the cultivated cucurbits, 101
White, Dr. David, 115
Woodson, Robert E., Jr.: Some dynamics of leaf variation in Asclepias tuberosa, 353
Wyoming, fossil plants from, 120

Y

Yeast, A new, 31
Yeasts: A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing mutants in, 85
Yeasts, see Saccharomyces

Z

Zea Mays, see Corn
Zea Mays, variation pattern of, 433
Zuñil, Guatemala, corn from, *440, 452, 464, 465*